

Does Recycling in Germinal Centers Exist?

Michael Meyer-Hermann

Institut für Theoretische Physik, TU Dresden, D-01062 Dresden, Germany
E-Mail: meyer-hermann@physik.tu-dresden.de

Abstract: A general criterion is formulated in order to decide if recycling of B-cells exists in GC reactions. The criterion is independent of the selection and affinity maturation process and based solely on total centroblast population arguments. An experimental test is proposed to verify whether the criterion is fulfilled.

1 Introduction

The affinity maturation process in germinal center (GC) reactions has been well characterized in the last decade. Despite a lot of progress concerning the morphology of GCs and the stages of the selection process, a fundamental question remains unsolved: Does recycling exist? *Recycling* means a back differentiation of antibody presenting centrocytes (which undergo the selection process and interact with antigen fragments or T-cells) to centroblasts (which do not present antibodies and therefore do not interact with antigen fragments but proliferate and mutate).

The existence of recycling in the GC has important consequences for the structure of the affinity maturation process in GC reactions. The centroblasts proliferate and mutate with high rates in the environment of follicular dendritic cells. During the GC reaction they differentiate to antibody presenting centrocytes which may then be selected by interaction with antigen fragments and T-cells. If the positively selected centrocytes recycle to (proliferating and mutating) centroblasts the antibody-optimization process in GCs may be compatible with random mutations of the centroblasts. On the contrary, one-pass GCs (without recycling) seem to be inconsistent with an optimization process which is based on undirected mutations. As a consequence, the search for mechanisms which facilitate the finding of the optimal antibody-type in GC reactions through somatic hypermutations is directly correlated to the existence of a centrocyte recycling process.

The recycling hypothesis was first formulated in 1993 [1] and there exist several approaches to verify it. (e.g. [2, 3, 4]). Experimentally, recycling of B-cells has never been observed directly. This may be due to a small number of recycled cells compared to the huge total number of cells in the GC. An indirect experimental evidence [2] for cell recycling was found by observing the rates of centrocyte apoptosis and the total cell population decrease after injection of a second antigen (different from the antigen which originally initiated the GC reaction) into a GC which already was in a later phase of its reaction. Both, apoptosis of centrocytes as well as the total cell population decrease were faster than in unperturbed GC reactions. Interestingly, the total cell population decrease was too fast to be solely explained by centrocyte apoptosis. This observation lead to the conclusion that a feedback mechanism from centrocytes to centroblasts should exist. Recycling is a good candidate for such a mechanism.

The major conceptual problem in the theoretical contributions is the number of assumptions which are necessary to describe the affinity maturation process. The recycling hypothesis was formulated because it seemed unlikely that a complex optimization process (involving up to about 9 somatic hyper-mutations [5, 6]) may occur in a one-pass GC, i.e. without recycling: Starting from about 12000 cells which mutate in a high-dimensional (shape) space the probability (mutation is likely to occur stochastically [7, 8]) for at least one cell to find the right mutant is very small. It would help, if the already affinity matured cells could reenter the proliferation process in order to multiply and to enhance the chance to find the optimal mutant.

Due to the origin of the recycling hypothesis, the efforts to show its existence were based on the influence of recycling on the maturation process. Consequently, it was found that the number of matured cells becomes substantially larger with recycling than without [3]. A quantitative description showed, that in order to bring the affinity maturation process into

accordance with experimental observations, the recycling probability of positively selected centrocytes should have as high values as 80% [4]. In both models important assumptions were made to represent the antibody-types in a shape space (affinity classes, dimension of shape space) and to allow a successive affinity maturation (affinity neighborhood).

Here, a new and simpler perspective is presented. Not the influence of recycling on the maturation process is examined but its influence on the total centroblast population. The centroblasts are believed not to present antibodies, so that they do not participate at the selection process via interaction with the antigens presented by the follicular dendritic cells. Therefore, in GCs without recycling the total centroblast population is independent of affinity maturation and selection. However, this statement may depend from the system under consideration. For instance, the centroblasts in GCs during splenic immune responses of mice has been observed to present antibodies [9].

The analysis starts from the presupposition that recycling does not exist. With a minimal set of assumptions (see Sec. 2) the implications for the time course of the total centroblast population are described in Sec. 3. The dependency of the results on some characteristic properties of various GCs is analyzed (see Sec. 3.1) and an experimental test of the implications is proposed (see Sec. 4).

2 Requirements

2.1 General assumptions

In this Gedanken experiment the total centroblast population $B(t)$ in a one-pass GC (without recycling) is considered. *Total* means, that the encoded antibody-type is ignored and all centroblasts enter the total population with equal weight (despite any differences in affinity to a presented antigen).

Presuming a one-pass GC no centrocytes are recycled to centroblasts, so that the total centroblast population becomes independent of the number of centrocytes. Basically, only centrocytes present antibodies and thus may undergo a selection process. As recoil effects to the total centroblast population are excluded, this magnitude is also independent of any properties or dynamics of the selection process. Especially, no influence of the already achieved affinity maturation during the optimization process exists. Also, neither the number of positively selected centrocytes, nor the speed of the selection process, nor the interaction with T-cells, nor the fate of the rejected centrocytes, nor the production of plasma and memory cells have any influence on the total centroblast population.

The total centroblast population changes exclusively according to centroblast proliferation and the differentiation of centroblasts to centrocytes. Therefore, $B(t)$ follows a differential equation of the following type

$$\frac{dB}{dt}(t) = -f(t)B(t) \equiv p(t)B(t) - g(t)B(t), \quad (1)$$

with time t , centroblast proliferation rate $p(t)$, rate of centroblast differentiation to centrocytes $g(t)$, and total rate function $f(t) = g(t) - p(t)$.

Cell proliferation has unquestionably to be described by an exponential increase of the population. Therefore, a linear proliferation term is used at every time of the GC reaction.

In the same way the differentiation term should be directly proportional to the centroblast population at every moment. Therefore, nonlinear terms do not enter the total centroblast population dynamics Eq. (1).

If the centroblasts participate at the selection process as observed in splenic immune responses in mice [9], i.e. if the centroblasts may, as the centrocytes do, undergo apoptosis, Eq. 1 remains correct but with a slightly modified interpretation: The growth of the number of centroblasts is inhibited by their apoptosis. Therefore, the function $p(t)$ denotes the proliferation rate of centroblasts reduced by the centroblast apoptosis rate. This should be beared in mind for the interpretation of the results.

Note, that no place is determined where proliferation or differentiation occurs. Therefore, the following argument is not restricted by the appearance or not-appearance of dark and light zones [10, 9] during a GC reaction.

2.2 About the rates

It is assumed that after an initial pure proliferation phase of some seeder cells, the total centroblast population does not further increase. This is in accordance with measurements of the follicle center volume, which show a peak 3 or 4 days after immunization [11]. As a consequence, the function $f(t)$ in Eq. (1) is positive during the whole GC reaction (started at $t = 0$), i.e. after the first pure proliferation phase.

The proliferation rate is probably constant during the whole GC reaction. This has not been shown experimentally in all details but is well established by a lot of independent data taken at different moments of the GC reaction [11, 12, 13]. All these data lead to the same large proliferation rate of

$$\frac{p(t)}{\ln(2)} \approx \frac{1}{6h}. \quad (2)$$

Nevertheless, for the following considerations the value of the proliferation rate will not be fixed. In order to assure generality of the argument proliferation rates according to Eq. (2) will be discussed as special case only. Therefore, the results of the following analysis apply also to GCs with a centroblast proliferation rate that varies in time.

The situation is more complicated in the case of the differentiation rate $g(t)$. This value is not known experimentally, such that its absolute number as well as its time course are unclear. From the injection of BrdUrd (5-bromo-2'-deoxyuridine) it is possible to identify the cells which were in cell cycle during the preceding 6 hours. In this way it was found in GCs with established dark and light zones, that the centrocyte population is renewed from centroblasts every 7 hours [11]. It follows for the differentiation rate that

$$\frac{g(t)}{\ln(2)} > \frac{1}{7h}. \quad (3)$$

Because of the lack of more detailed information, different possibilities for the time dependence of the centroblasts to centrocytes differentiation rate will be considered in the following argument.

2.3 A standard germinal center

Before analyzing the implications of this Gedanken experiment a typical GC has to be defined as reference system. The GC reaction is initiated by a few seeder cells in the environment of follicular dendritic cells (FDC) [5, 11, 14, 15]. These cells proliferate for about 3 days, and give rise to about 12000 centroblasts [11]. At this time $t_0 = 0h$ the intrinsic GC reaction starts, including a continuation of proliferation and, additionally, the differentiation of centroblasts to centrocytes. The analysis made here does not consider the first phase of pure proliferation, but deals solely with the GC properties in the second phase. The life time T of a germinal center is about 18 days ($T = 432h$) [11, 16], in which the first phase of pure proliferation is not included. At the end of the GC reaction, only a few cells remain in the FDC environment [11, 16], which is assumed to be about 5 cells in a standard GC. The sensitivity of the results on these assumptions will be analyzed (see Sec. 3.1).

3 Analysis of the centroblast population

The different time courses considered for the total centroblast population of a standard GC (defined in Sec. 2.3) are shown in Fig. 1. An exponential decrease corresponds to a constant rate function (see Fig. 2 dashed line). A linear population decrease is achieved with the rate function Eq. (15) (see Fig. 2 dotted line) which is derived in the Appendix. Finally, using a quadratic rate function Eq. (17) (see Fig. 2 full line) the centroblast population remains unchanged for the major duration of the GC reaction and is steeply reduced at the very end.

Experimentally, the time course of the GC total centroblast population seems to lie in between the exponential and the linear scenario [11, 17]. Anyhow, the scenario with a long-lasting constant centroblast population, i.e. with the quadratic rate function, is certainly a reasonable upper bound for possible time courses of the total centroblast population (Fig. 1 full line). On the other hand, a population decrease that is faster than exponential is unlikely.

It follows that during the major life time of GCs ($t < 15d$) the condition

$$f(t) = g(t) - p(t) < 0.018/h \quad (4)$$

holds (see Fig. 2). On the other hand the requirement of a monotonically decreasing centroblast population leads to the bound $f(t) > 0$, i.e. $g(t) > p(t)$. Taking these results together it follows

$$p(t) < g(t) < p(t) + 0.018/h, \quad (5)$$

which is a very powerful condition. This means that in an experiment the centroblast to centrocyte differentiation rate $g(t)$ has to fulfill this condition during the first 15 days of the GC reaction. If it turns out that the above condition is not guaranteed experimentally, then – assuming the rather weak requirements made to be valid – a GC reaction without recycling of centrocytes to centroblasts is ruled out.

Note, that assuming the proliferation rate to adopt the value Eq. (2), the above condition can be reformulated. It follows a lower bound for the differentiation rate which is in




Figure 1: Considered time courses of the centroblast population during GC reactions. The model results remain valid for all GCs with centroblast populations within these extreme scenarios.

accordance with Eq. (3):

$$\frac{1}{6h} < \frac{g(t)}{\ln(2)} < \frac{1}{5.19h} . \quad (6)$$

This means that if GCs without recycling exist, a centroblast shall (in average) take not less than 5.19 hours to differentiate into centrocytes.

Supposing that the linear centroblast population decrease is the realistic scenario, the upper bound becomes even more powerful. During the first 10 days the condition

$$p(t) < g(t) < p(t) + 0.005/h \quad (7)$$

must hold in a GC reaction without recycling. With the proliferation rate in Eq. (2) this corresponds to an average centroblast differentiation time of at least 5.75 hours. This result demonstrates, that the condition for a one-pass GC may become stronger than in Eqs. (5) and (6), depending on the characteristics of the total centroblast population decrease.




Figure 2: The function $f(t) = g(t) - p(t)$ corresponding to the different time courses of the centroblast population in Fig. 1. The GC life time is $T = 432 h$. During the major life time of the GC the rate function respects the upper bound $f(t) < 0.018/h$ (see Eq. 4).

3.1 Variation of the GC properties

3.1.1 The case of constant rates

In order to check the robustness of the results Eqs. (5) and (6) the most critical scenario (i.e. with weakest conditions) is considered in some more detail: the scenario with constant rates. If the proliferation and differentiation rate are constant, the centroblast population model reduces to a pure exponential decrease of the population

$$B(t) = B(0) \exp((p - g)t), \quad (8)$$

where the constant rate g is unknown. Focusing on the population after $T = 18$ days of the GC reaction, the rate function

$$f = g - p = -\frac{1}{T} \ln \left(\frac{B(T)}{B(0)} \right) \quad (9)$$

is found. The resulting values for the differentiation rate are shown in Fig. 3 in dependence of the GC life time and for different numbers of remaining cells at the end of the reaction.

As expected the rate becomes larger for shorter GC life times T . Even supposing a rather




Figure 3: The values of the (constant) rate function f in the case of an exponentially decreasing centroblast population. The dependence of condition Eq. 5 of the GC life time and of the number of remaining centroblasts at the end of the GC reaction is shown to be weak (see also Eq. 10 for the most pessimistic case).

short life time of 12 days and further supposing that only one cell remains at the end of the GC reaction, one gets an upper bound of

$$p(t) < g(t) < p(t) + 0.033/h \quad (10)$$

for possible values of the differentiation rate. Incorporating the proliferation rate Eq. (2) it follows

$$\frac{1}{6h} < \frac{g(t)}{\ln(2)} < \frac{1}{4.7h}. \quad (11)$$

As a consequence, the statement that in one-pass GCs (without recycling) the centroblast to centrocyte differentiation rate should respect the above bounds Eqs. (5) and (6) is not altered dramatically by variation of the GC properties.

The initial number of centroblasts $B(0)$ was not varied so far. Note that only the ratio $B(T)/B(0)$ enters Eq. (9), so that a variation of the initial number of centroblast is

equivalent to the variation of the final number of remaining cells $B(T)$. Therefore, the constraints Eqs. (10) and (11) are also valid with e.g. 60000 initial and 5 remaining cells. Lower numbers of initial cells allow even stronger bounds than the above ones.

3.1.2 The case of linear centroblast decrease




Figure 4: The time course of the rate function $f = g - p$ (corresponding to a linear centroblast population decrease) for different GC life times T and for different numbers of remaining centroblasts $B(T)$ at the end of the GC reaction. Condition Eq. 7 turns out not to depend on the number of remaining centroblasts. The rate function is sensitive to the total GC life time in the late phase of the reaction. However, condition Eq. 5 remains valid for practically the whole GC life time.

In the case of a linear centroblast population decrease (see Fig. 1 pointed line) the corresponding rate function $f(t)$ (see Eq. (15)) is analyzed for various GC life times T and numbers of remaining blasts at the end of the reaction $B(T)$. The result is shown in Fig. 4. Condition Eq. (7) is not affected at all by a variation of the number of remaining cells at the end of the GC reaction in a range of three orders of magnitude (the values chosen for $B(T)$ are 0.1, 5, and 100). The three corresponding curves are practically not distinguishable (see Fig. 4).

The situation is different for the variation of the GC life time T . Here, the period in which condition Eq. (7) remains valid is prolonged for long living GCs and shortened for short living GCs (see Fig. 4 dashed and pointed line, respectively). For a very short life time of 12 days (plus 3 days of pure proliferation) the condition remains valid for 4 days only (7 days after immunization). However, condition Eq. (5) remains valid even in this case for 11 days, so for approximately the whole GC life time.

4 Conclusions

The above analysis is summarized with the statement that in a one-pass GC (without recycling) the centroblast to centrocyte differentiation rate should respect the strong condition Eq. (5) $p(t) < g(t) < p(t) + 0.018/h$ during the first 15 days of the GC reaction (with standard properties which are defined in 2.3). For a standard proliferation rate Eq. (2) of $\ln(2)/6h$ this translates into an average time for centroblasts to differentiate into centrocytes which cannot be substantially shorter than 5 hours. If this condition is not fulfilled, recycling has necessarily to be present in GC reactions. Note that, the other way round, if the average differentiation of centroblasts to centrocytes takes more than 5 hours, recycling is not necessarily absent.

This statement is based essentially on two (weak) assumptions: Firstly, linear population dynamics (see Eq. (1)) for the total centroblast population in GCs. Secondly, a monotonic decrease of the total centroblast population. The conclusions apply universally to GCs with various characteristics due to the independence of all speculations on shape spaces, selection processes, and affinity maturation. However, one should be aware that still some assumptions may be questioned. Remind, for instance, that for GCs with selectable centroblasts (that may undergo apoptosis) the rate p has to be interpreted as the rate of centroblast proliferation reduced by the centroblast apoptosis rate.

In order to decide if recycling does exist in GC reactions one has to determine the centroblast to centrocyte differentiation rate experimentally *at least* at one representative moment of the GC reaction. The moment has to be chosen such that on one hand the GC reaction is already well established. On the other hand, the later the measurement is done the less is known on the proliferation rate. In addition, one approaches the regime where the above condition loses its validity (see Fig. 2). Therefore, an optimal time will be about 8 days after immunization. However, if one expects the centroblast proliferation rate to vary substantially during this central phase of GC reaction, the standard values for p cannot be used. In this case, one would have to determine not only the centroblast differentiation rate but also the centroblast proliferation rate.

Technically, the centroblast differentiation rate may be measured by labeling studies of B-cells in the GC which are in cell cycle as it was done in [9, 11].

Alternatively, a systematic and time resolved analysis of the total centroblast population at different moments of the GC reaction may improve the criterion (see e.g. Eq. (7)) because one could distinguish which of the cell population scenarios presented in Fig. 1 corresponds to real GC reactions. This may be done by the measurement of the GC volume at several times of the GC reaction as in [11, 17, 18].

Appendix

The differential equation Eq. (1) is formally solved by

$$B(t) = B(t_0 = 0 \text{ h}) \exp \left(- \int_0^t dx f(x) \right) . \quad (12)$$

In the following, the border conditions (life time of GC T , and centroblast population at $t_0 = 0 \text{ h}$ and T) are incorporated into the rate function f .

To achieve a linear decrease of the centroblast population, the second time derivative of the population is required to vanish. From Eq. (1) and using Eq. (12) it follows

$$\begin{aligned} \frac{d^2 B}{dt^2} &= -\frac{df}{dt} B - f \frac{dB}{dt} \\ &= -\frac{df}{dt} B(0) \exp \left(- \int_0^t dx f(x) \right) + f B(0) f \exp \left(- \int_0^t dx f(x) \right) . \end{aligned} \quad (13)$$

This expression vanishes for

$$\frac{df}{dt} = f^2 , \quad (14)$$

which is solved by

$$f(t) = -\frac{1}{t - c_1} . \quad (15)$$

The integration constant c_1 is fixed by the condition that the final population at time T is given by $B(T)$. With Eq. (12) this leads to

$$c_1 = \frac{T}{1 - \frac{B(T)}{B(0)}} , \quad (16)$$

which is a positive constant for $B(T) < B(0)$.

In the case of a quadratic rate function

$$f(t) = \frac{(1 \text{ h})}{(t - c_2)^2} \quad (17)$$

an analogous calculation which incorporates the final centroblast populations leads to

$$c_2 = \frac{T}{2} + \sqrt{\frac{T^2}{4} - \frac{(1 \text{ h}) T}{\ln \left(\frac{B(T)}{B(0)} \right)}} . \quad (18)$$

Other solutions are formally possible but do not lead to biologically reasonable centroblast populations.

References

- [1] Kepler, T.B., and Perelson, A.S., Cyclic re-entry of germinal center B cells and the efficiency of affinity maturation. *Immunol. Today* **14** (1993), 412-415.

- [2] Han, S.H., Zheng, B., Dal Porto, J., Kelsoe, G., In situ Studies of the Primary Immune Response to (4-Hydroxy-3-Nitrophenyl) Acetyl IV. Affinity-Dependent, Antigen-Driven B-Cell Apoptosis in Germinal Centers as a Mechanism for Maintaining Self-Tolerance. *J. Exp. Med.* **182** (1995), 1635-1644.
- [3] Oprea, M., Nimwegen, E. van, and Perelson, A.S., Dynamics of One-pass Germinal Center Models: Implications for Affinity Maturation. *Bull. Math. Biol.* **62** (2000), 121-153.
- [4] Meyer-Hermann, M., Deutsch, A., and Or-Guil, M., Recycling probability and dynamical properties of germinal center reactions. *J. Theor. Biol.* **210** (2001), 265-285.
- [5] Küppers, R., Zhao, M., Hansmann, M.L., and Rajewsky, K., Tracing B Cell Development in Human Germinal Centers by Molecular Analysis of Single Cells Picked from Histological Sections. *EMBO J.* **12** (1993), 4955-4967.
- [6] Wedemayer, G.J., Patten, P.A., Wang, L.H., Schultz, P.G., and Stevens, R.C., Structural insights into the evolution of an antibody combining site. *Science* **276** (1997), 1665-1669.
- [7] Weigert, M., Cesari, I., Yonkovitch, S., and Cohn, M., Variability in the light chain sequences of mouse antibody. *Nature* **228** (1970), 1045-1047.
- [8] Radmacher, M.D., Kelsoe, G., and Kepler, T.B., Predicted and Inferred Waiting-Times for Key Mutations in the Germinal Center Reaction - Evidence for Stochasticity in Selection. *Immunol. Cell Biol.* **76** (1998), 373-381.
- [9] Camacho, S.A., Koscovilbois, M.H., and Berek, C., The Dynamic Structure of the Germinal Center. *Immunol. Today* **19** (1998), 511-514.
- [10] Nossal, G., The molecular and cellular basis of affinity maturation in the antibody response. *Cell* **68** (1991), 1-2.
- [11] Liu, Y.J., Zhang, J., Lane, P.J., Chan, E.Y., and MacLennan, I.C.M., Sites of specific B cell activation in primary and secondary responses to T cell-dependent and T cell-independent antigens. *Eur. J. Immunol.* **21** (1991), 2951-2962.
- [12] Hanna, M.G., An autoradiographic study of the germinal center in spleen white pulp during early intervals of the immune response. *Lab. Invest.* **13** (1964), 95-104.
- [13] Zhang, J., MacLennan, I.C.M., Liu, Y.J., and Land, P.J.L., Is rapid proliferation in B centroblasts linked to somatic mutation in memory B cell clones. *Immunol. Lett.* **18** (1988), 297-299.
- [14] Jacob, J., Kassir, R., and Kelsoe, G., In situ studies of the primary immune response to (4-hydroxy-3-nitrophenyl)acetyl. I. The architecture and dynamics of responding cell populations. *J. Exp. Med.* **173** (1991), 1165-1175.

- [15] Kroese, F.G., Wubbena, A.S., Seijen, H.G., and Nieuwenhuis, P., Germinal centers develop oligoclonally. *Eur. J. Immunol.* **17** (1987), 1069-1072.
- [16] Kelsoe, G., The germinal center: a crucible for lymphocyte selection. *Semin. Immunol.* **8** (1996), 179-184.
- [17] Hollowood, K., and Macartney, J., Cell kinetics of the germinal center reaction – a stathmokinetic study. *J. Immunol.* **22** (1992), 261-266.
- [18] de Vinuesa, C.G., Cook, M.C., Ball, J., Drew, M., Sunners, Y., Cascalho, M., Wabl, M., Klaus, G.G.B., and MacLennan, C.M., Germinal Centers without T Cells. *J. Exp. Med.* **191** (2000), 485-493.

Acknowledgments

I thank Andreas Deutsch and Michal Or-Guil for intense discussions and valuable comments.